

Xenon Recovery from Used Nuclear Fuel

For Tests of Fundamental Physics

Regan Ross, June 25, 2026

CAP Congress



McGill
UNIVERSITY

The Standard Model

The SM has been extremely successful in describing particles and their interactions—even predicting the existence of new particles.


However, we know it is incomplete.



Baryon Asymmetry

Physics is symmetric under the interchange of particle and anti-particle

But we see only *matter*, not *antimatter* in the universe.



Dag und Nacht, M.C. Escher, 1938

Dark Matter


There is abundant evidence for celestial mass of an unknown nature:

Conflicting predicted and measured galactic rotation

Gravitational lensing (strong and weak)

Galaxy clustering dynamics

Cosmology and large scale structure formation in the universe



Still life Reslicitando, R. Varo, 1963

Neutrino Masses

Neutrino masses are much smaller than other particle masses

The SM has little to say about their mass generation mechanism, their oscillatory nature, or their absolute mass scale



Le chant d'amour, R. Magritte, c. 1962

Neutrino Masses

Neutrino masses are much smaller than other particle masses

The SM has little to say about their mass generation mechanism, their oscillatory nature, or their absolute mass scale

Or their mass ordering



Ceci n'est pas un neutrino, R. Magritte, c. 2026 (AI)

*My advice is to go for the messes
- that's where the action is.*

Steven Weinberg, McGill Science Convocation, 2003

The “Rare Event Search”

Searches for two kinds of hypothesized interactions:


Dark matter - nuclear interaction

- DM scatters with nucleus
- Produces detectable signal
- Would establish particle nature of DM
- Establishes DM mass scale




Neutrinoless Double Beta Decay

- Electrons emitted with equal and opposite momentum
- *Lepton number* not conserved
- Establishes neutrino mass scale



The Nucleus as a Laboratory


- Undergoes double beta decay
- Heavy nuclei are good for DM detection
- Strong scintillation and ionization signals possible
- Can be liquefied at $\sim 165\text{K}$
- Chemically inert
- *Comparatively* high natural abundance



Xenon Time Projection Chambers

For $0\nu\beta\beta$ and Dark Matter Searches

- Xenon time projection chambers (TPCs) have been continuously improving for over 2 decades
- They leverage xenon's favourable properties to detect light, charge, event position and topology.
- Event parameters and self-shielding provides background discrimination



D. Gallacher

$0\nu\beta\beta$ Candidate Isotopes

should meet the following criteria:

- High natural abundance; large N
- Fine energy resolution; small ΔE
- Few intrinsic backgrounds; small B
- Easily discernible $0\nu\beta\beta$ signal
- Feasible to work with

$$T_{1/2}^{0\nu} \leq \frac{\epsilon \ln 2 \sqrt{Nt}}{\sqrt{B\Delta E}}$$

$0\nu\beta\beta$ Candidate Isotopes

should meet the following criteria:

$$T_{1/2}^{0\nu} \propto \sqrt{Nt}$$


Xenon-Based Experiments

For $0\nu\beta\beta$ and Dark Matter Searches

There is evident increasing demand for xenon for WIMP dark matter search experiments and $0\nu\beta\beta$ searches.


Xenon is becoming the single greatest expenditure!

At $\approx$ \$5/g, tonne scale experiments can cost \$5M to \$50M / tonne !




Atmospheric Abundance of Xe

Dilute, but unlimited.





Fission of ^{235}U



CANDU Fission Products

CANDU = CANadian Deuterium Uranium




Burnup of 685 GJ/kg U, One year cooled.

Xenon from Used Nuclear Fuel

An untapped source

Bubbles of Kr and Xenon are trapped in the UO_2 fuel after burn-up. They accumulate at grain boundaries in the metal.

~ 1.5 g of Xenon per kg of U metal



How much is there?

An untapped source

3.3 million used CANDU fuel bundles


64.260 kt of heavy metal (t-HM)

80 t of xenon → 36 t of Xe-136

Inventory will **double** by reactor expiration.

Xenon-136 present at about 560 ppm.

14,000 times more concentrated
than atmospheric xenon.




What else is there?

Isotope	Bq/kg U *	$T_{1/2}$	Mode	E (keV)	g/kg U *
^3H	2.96e+09	12.32 y	β^-	19	8.27e-06
^{81}Kr	4.26e+00	2.29e+05 y	EC	—	5.45e-09
^{85}Kr	4.55e+10	10.76 y	β^-	687	3.13e-03
^{129m}Xe	0.00e+00	8.9 d	IT	236	3.51e-07
^{131m}Xe	0.00e+00	11.8 d	IT	164	0.00
^{133}Xe	0.00e+00	5.25 d	β^- / γ	427 / 81	0.00
^{219}Rn	1.04e+01	3.96 s	α	6946	2.15e-20
^{220}Rn	2.70e+04	55.6 s	α	6405	7.92e-16
^{222}Rn	2.18e+00	3.82 d	α	5590	3.82e-16

* after 10 years of cooldown

Potential Background Sources



Demonstrable Purification Ability

- Kr Removal with Cryogenic Distillation, XENON Collaboration, [arXiv:1612.04284](#)
 - Down to < 30 ppq
- Rn Removal down to Solar ν Level, XENONnT Collaboration [arXiv:2502.04209](#)
 - Down to $< 1\mu\text{Bq} / \text{kg}$
- Rn Removal (from Ar) with Silver Zeolites, Dunford et al., [arXiv:2505.07979](#)
 - Using novel adsorbents

Operational Challenges

Neglecting bureaucratic friction

- Maximum $\sim 1\text{g Xe}$ / element (37 per bundle)
- Fuel elements (pellets) are designed to HOLD gases
- Dangerous (dose rates of 10s to 100s of Sv/hr)
- Experimentation must be done in hot cells



(Pre-irradiation ;)

Operational Challenges

- Hot cells are expensive ($\sim \$1000/\text{hr}$)
- Instruments in hot cells must be operated by “manipulators”
- Apparatus must be built accordingly




Hot cell at CNL

Operational Challenges



Hot cell at CNL



Hot cell at Nevada Test Site

Operational Challenges



UNF Transportation Cask (DOE)

A CNL Visit

(no photos allowed)

- Extensive facilities for analysis & fuel QA
- Focus on reactor engineering
(as opposed to basic research)
- Not opposed to collaboration
- Do not have a supply of UNF



CNL on the Ottawa River

Conclusion

Where do we go from here?

Remarks

- The most plausible route is to build a parasitic process on top of *reprocessing*
- Maybe this is worth exploring without the burden of creating a new supply, but merely promoting new science & collaboration

Questions

- Can we construct a lab-scale demonstrator?
Is it worth trying?
- Who can help?
- Can this lead to collaboration with La Hague?
- Can this grow to support tonne-scale experiments?

Thank you!





Government of Canada
Gouvernement du Canada

MENU

[Canada.ca](#)

Government of Canada launches Nuclear Energy Strategy

From: [Natural Resources Canada](#)

News release

June 22, 2026

Newmarket, Ontario

(Monday)

Contact:
Regan Ross, regan.ross@mail.mcgill.ca




McGill
UNIVERSITY


PUREX Process

The current standard

- UNF fully dissolved in acid
- Metals precipitated out (U, Pu)
- Gases are generally vented (all of them.)
- Could they be captured and purified?



UNF storage at La Hague, France.



XLZD

The definitive rare event observatory

- Designed to reach world-leading sensitivity down to the neutrino fog in the WIMP search
- Anticipated sensitivity to $0\nu\beta\beta$ close to 10^{28} y
- International collaboration with 450+ scientists
- Could be attracted to SNOLAB 
- Goal of a final detector with 60-80 T of liquid Xe in a T
- Could be upgraded with enriched xenon

